

ORIGINAL ARTICLE

Title: Virtual Reality for Disaster Response Training: Feasibility and Impact on
Emergency Healthcare Professionals

Authors: Edmund Wong Kwang Yuen, Mohd Hisham Isa, Ismail Mohd Saiboon, Mohd
Johar Jaafar, Ida Zarina Zaini

Submitted Date: 10-03-2025

Accepted Date: 16-09-2025

Please cite this article as: Edmund Wong Kwang Yuen, Mohd Hisham Isa, Ismail Mohd Saiboon, Mohd Johar Jaafar, Ida Zarina Zaini. Virtual Reality for Disaster Response Training: Feasibility and Impact on Emergency Healthcare Professionals. Education in Medicine Journal. (early view).

This is a provisional PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article.

Virtual Reality for Disaster Response Training: Feasibility and Impact on Emergency Healthcare Professionals

Edmund Wong Kwang Yuen, Mohd Hisham Isa, Ismail Mohd Saiboon, Mohd Johar Jaafar, Ida Zarina Zaini

Department of Emergency Medicine, Faculty of Medicine, Universiti Kebangsaan Malaysia, Cheras
Kuala Lumpur, Malaysia.

Corresponding Author

Mohd Hisham Isa, Department of Emergency Medicine, Faculty of Medicine UKM, 56000 Cheras Kuala Lumpur, Malaysia

Email: mohimois@ukm.edu.my

ABSTRACT

Disaster medicine training is essential for healthcare professionals and incorporates various modalities, including classroom-based education, tabletop exercises, drills, and field exercises. Advancements in technology, such as the development of Extended Reality (XR) modalities, have led to the emergence of Virtual Reality (VR) as an effective educational tool. This is particularly relevant with regard to Disaster Response Medicine (DRM), enhancing experiential learning and preparedness. Accordingly, this study evaluates the feasibility of integrating VR into DRM training. As such, a cross-sectional study was conducted among emergency department doctors. Participants engaged in a VR-based scenario for emergency response training and provided feedback through a 5-point Likert scale questionnaire assessing usability, personal experience, and adverse events. Furthermore, a quantitative performance matrix was employed to evaluate participant performance levels. A total of 70 participants volunteered for the study, with 38 (54.3%) being male and 32 (45.7%) being female, with a median age of 33. Notably, 49 participants (70%) had no prior experience with VR. Over 80% reported the VR system was easy to use, relevant to real-life scenarios, and beneficial to their field, while more than 90% believed it could enhance professional performance. Meanwhile, the most common adverse event was visual discomfort (30%), followed by dizziness, with fewer reports of headaches, fatigue, nausea, stomach awareness, and postural instability. Despite these effects, 60% of participants exceeded the 80% performance benchmark. Moreover, VR-based training in DRM demonstrated high usability, realism, and effectiveness in skill enhancement. Nevertheless,

with optimized exposure time and user-friendly interfaces, VR can serve as a viable and immersive tool to complement traditional DRM training while minimizing adverse effects.

Keywords: Disaster response medicine, virtual reality, healthcare training, emergency medicine training, simulation-based learning.

INTRODUCTION

Disasters are unpredictable and infrequent events, making real-world exposure to such scenarios limited in everyday clinical practice (1). Nonetheless, for healthcare professionals, particularly those in the emergency department, being prepared to respond effectively to disasters is critical (2). Thus, comprehensive disaster education and training are essential to equip medical personnel with the necessary knowledge, decision-making skills, and hands-on proficiency required in high-pressure, mass-casualty situations (3).

Traditional training methods, such as classroom-based learning, tabletop exercises, and field training exercises, provide valuable theoretical and practical insights. However, they may not always replicate the dynamic, high-stakes environment of real disasters. According to Hsu et al. (4), traditional classroom teaching and web-based training often lack realism and excitement, leading to varying levels of engagement among participants. Furthermore, organizing these exercises presents significant challenges, as they require extensive preparation, resource allocation, logistical coordination, and the involvement of a large number of participants (5-6). In addition, it should be noted that the implementation of real-life drills can have the unintended consequence of disrupting regular local services. This, in turn, poses additional obstacles in terms of coordinating training efforts (7).

Innovative training approaches such as Virtual Reality (VR) provide an immersive and interactive platform that bridges the gap between theoretical knowledge and practical application in disaster response preparedness (7-8). In line with this, VR offers unique advantages in education and training by enabling realistic simulations, customizable scenarios, and performance tracking (9). Additionally, the utilization of VR as a learning strategy has been observed to significantly enhance student engagement, attention, and interaction, resulting in a beneficial impact (10). Andonova et al. (11) demonstrated that VR significantly improves information retention and recall by engaging multiple senses, creating more memorable learning experiences. In essence, VR supports personalized learning, allowing users to train at their own pace on any modern laptop or desktop without the need for specialized software.

The vast potential of VR in disaster response training is particularly noteworthy. Therefore, this study examines the feasibility of using a locally developed VR module to train emergency healthcare

professionals in Disaster Response Medicine (DRM). Specifically, the assessment focuses on usability, user experience, potential adverse events, and performance outcomes.

METHODOLOGY

Study Design and Participants

A cross-sectional study was conducted in the VR facility at the Faculty of Medicine between May 2022 and December 2023. This study was approved by the Medical Research and Ethics Committee (JEP2022-333). Participants were emergency doctors with varying levels of experience in clinical practice from a single university hospital's emergency department in Klang Valley. Consent was obtained from all participants before the study commenced. Individuals with a history of significant motion sickness, anxiety disorders, or epilepsy were excluded. A convenience sampling method was employed to recruit participants.

Development of a Disaster Response VR Module

A disaster response VR module was developed in collaboration with three emergency physician consultants, a healthcare simulation specialist, and a VR technology team. The module simulates a mass casualty incident involving a bus crash into a ravine, requiring participants to resume the role of prehospital care provider to manage the scene. Key tasks include reporting to the incident commander, organizing a Casualty Clearing Station (CCS), placing medical resources, and constructing a medical facility. Participants use a VR controller to navigate the scenario, respond to embedded multiple-choice questions, and make critical decisions. They are required to establish a CCS by selecting an appropriate location and choosing the necessary equipment using the controller. These activities assess participants' knowledge, decision-making, and clinical reasoning skills. The scenario includes 18 virtual patients with variable injuries, triaged into four categories: critical (red), semicritical (yellow), non-critical (green), and deceased (black). Treatment is administered at the CCS prior to hospital transport. During each segment of the disaster response VR simulation, participants were required to attempt the questions presented. Upon selecting the correct answer, the VR system provided explanations to enhance self-directed learning (Figure 1). Content and construct validation of the VR disaster response training module were conducted by two senior consultant emergency physicians with expertise in disaster response management. They reviewed and evaluated the module, offering feedback on its technical aspects. The research team subsequently refined the module based on

their recommendations. Face validity was assessed by a group of ten medical officers who were not involved in the study.




Figure 1: VR Simulation Interface

Figure 1: VR simulation interface displaying multiple-choice questions with feedback

Study Protocol

Participants were assigned to small groups, and they were provided with a 30-minute instructional video covering key principles of DRM. Prior to the intervention, an orientation session familiarized the participants with the VR controller, VR Stereo 3D Glass (Figure 2), and the appropriate positioning for the duration of the VR study was conducted. The intervention consisted of a single session of immersive VR simulation training. Participants were equipped with VR headsets and interacted with the virtual environment through motion-tracked controllers, allowing them to select the correct answers from the options displayed on the screen. They were assessed across five specific areas: i. Scene assessment and team reporting for duty, ii. Selection of an appropriate CCS, iii. Triage based on injury severity, iv. Selection of appropriate treatments, and v. Prioritization of victim transport based on injury severity. To ensure the safety of participants, the duration of the VR simulation was limited to a maximum of 30 minutes. Participants were continuously monitored to prevent accidental collisions with surrounding objects. They

were given the option to pause the VR simulation if they experienced any undesirable outcomes. The entire session was recorded using a Sony ZV-1 camera, and performance data were analyzed using an unbiased evaluation matrix. Upon completing the module, participants completed the questionnaire for further analysis.



Figure 2: VR Stereo 3D Glasses and Controllers

Figure 2: Virtual Reality Stereo 3D Glasses and Controllers

Assessment tools

Questionnaire

The questionnaire assessed participants' socio-demographic characteristics, usability, and experience with VR simulation training in DRM, and any adverse events encountered. Usability and user experience were evaluated using a 5-point Likert scale, ranging from 1 (strongly disagree) to 5 (strongly agree). Responses of "Strongly Agree" and "Agree" were categorized as positive, while "Disagree" and "Strongly Disagree" were classified as negative. "Neutral" was considered an absence of strong preference. Moreover, to

evaluate adverse events associated with VR training, a rating scale was used to assess symptoms such as headaches, dizziness, visual discomfort, fatigue, nausea, vomiting, and postural instability. Responses were scored from -2 (strongly disagree) to +2 (strongly agree), with lower (more negative) scores indicating a lower occurrence of side effects, meaning participants experienced fewer adverse events. The questionnaire was validated through a two-step process. Face validation was performed with feedback from 10 independent Emergency Department medical officers, while content validation was conducted by six experienced emergency physicians from different institutions. They evaluated and scored the questionnaire to ensure it effectively measured the intended constructs. The Face Validation Index (FVI) and Content Validation Index (CVI) were calculated, yielding high agreement scores of 0.97 and 0.98, respectively.

Objective Performance Matrix

Participants' performance was assessed using an objective scoring system. They were required to demonstrate key competencies. This includes reporting to the incident commander, identifying their team, assessing the situation, selecting an appropriate location for CCS, establishing a CCS by accurately placing medical icons, and conducting triage evaluations. Additionally, they were evaluated on their selection of appropriate treatment and transportation options. Each correct action was awarded one point, with a maximum possible score of 62. Scores were then converted into percentages and categorized into three performance levels: high ($\geq 80\%$), moderate (61% to 79%), and low ($< 60\%$). The objective performance matrix was validated by two emergency physicians with expertise in disaster management from an independent institution. Face validity was further assessed by a group of medical officers.

Statistical analysis

All statistical analyses were conducted using the Statistical Package for the Social Sciences (SPSS) version 27.0. Descriptive statistics were applied to summarize all data, including demographic data, Likert scale questionnaire responses, and a performance matrix. Notably, age was reported as the median with Interquartile Ranges (IQR), while categorical demographic variables were presented as frequencies and percentages. Adverse events were assessed using a rating scale ranging from -2 ("strongly disagree") to +2 ("strongly agree"), where lower scores indicated a lower likelihood of experiencing a specific adverse effect. Performance assessments were categorized into three levels based on Bloom's cutoff points: high, moderate, and low. The results were summarized using frequency and percentage distributions.

RESULTS

Demographic: A total of 70 participants were involved in the study. Among them, 38 (54.3%) were male, and 32 (45.7%) were female. The median age of the participants was 33 years. Notably, 49 participants (70%) had no prior experience with VR. Similarly, 69 participants (98.6%) had never used VR simulation for disaster training (Table 1).

Table 1: Sociodemographic characteristics of the participants (n= 70)

Variable	N (%)
Age	33 (32,36)*
Gender	
Male	38 (54.3)
Female	32 (45.7)
Wearing Glasses	
Yes	40 (57.1)
No	30 (42.9)
VR Games Experience	
Yes	21 (30.0)
No	49 (70.0)
VR Simulation Experience in Disaster Training	
Yes	1 (1.4)
No	69 (98.6)

Note: * Median (IQR)

Usability of VR in DRM Training

Figure 3 presents participants' responses regarding the usability of VR in DRM training. The majority (81.4%) agreed that the VR simulation user interface was user-friendly. Most participants (92.9%) perceived the VR scenarios as realistic and potentially occurring in real life. Moreover, 91.4% indicated they would act in a similar manner during a real-life situation. In contrast, the visual portrayal received

relatively lower positive feedback, with 64.3% agreeing it was realistic, 21.4% responding neutrally, and 14.3% expressing negative perceptions.




Figure 3: Response of participants on Usability of VR in Training DRM

Figure 3: Response of participants on usability of VR in training DRM

VR Experience in DRM Training

Figure 4 summarizes participants' responses regarding their VR experience in DRM training. A substantial proportion agreed that VR simulation was effective (80.0%) and realistic (71.4%) in DRM teaching. More than half (55.7%) believed VR addressed training needs not covered by the usual curriculum, though 32.9% were neutral. Most participants reported VR to be applicable to their practice (84.3%) and helpful for improving it (92.9%). Enjoyment of VR in disaster training was high (92.9%), with 90.0% recommending its use in future training. Additionally, 71.4% perceived VR as more cost-effective than traditional methods.




Figure 4: Participants' Responses on VR Experience in DRM

Figure 4: Participants' responses on VR experience in DRM training

Adverse Events in DRM Training Using VR

Among the reported symptoms, visual discomfort was the most common, affecting 30.0% of participants. Dizziness or spinning sensations were reported by 14.3%, while fatigue was noted by 8.6%. Headache and postural instability were each experienced by 7.1% of participants, and nausea-related symptoms, including vomiting and stomach discomfort, were the least frequently reported (2.9%). The scoring system reflected this trend, with more negative values corresponding to lower symptom frequency. Nausea-related symptoms had the lowest score (-99), followed by postural instability (-89) and headache (-73) (Table 2).

Table 2: Adverse Events using VR in DRM Training

Variables	Positive symptoms n (%)	Scoring system
I experienced headache during VR training	5 (7.1)	-73
I had dizziness/spinning sensation during VR training	10 (14.3)	-58
I had visual discomfort during VR training	21 (30)	-29
I had fatigue and tiredness during VR training	6 (8.6)	-71
I had nausea/vomiting/stomach awareness due to VR training	2 (2.9)	-99
I had postural instability during VR training	5 (7.1)	-89

VR Performance in DRM Training

The performance level among participants in the VR DRM training is presented in Figure 5. The majority of participants (60%, n = 42) achieved a high level of performance, while 40% (n = 28) demonstrated moderate performance. Notably, no participants were categorized as having a low level of performance.




Figure 5: VR Performance in DRM Training

DISCUSSION

This study evaluated the feasibility of VR-based training in DRM and demonstrated promising results. The findings indicate that VR simulation was well received by participants, with high usability ratings and a strong perception of realism. The effectiveness of usability was attributed to the precise replication of disaster situations that aligned with the educational requirements of the participants. VR offers an unparalleled degree of realism and immersion (12), enabling trainees to interact with a realistic environment that nearly replicates actual disaster scenarios. Additionally, this VR system offered feedback mechanisms to help users discover areas for growth, implement necessary changes, and continuously better their skills. Numerous studies highlight the crucial role of feedback in guiding student progress and optimizing performance (13-14). This positive reaction emphasizes the need for having an instinctive and easily understandable interface for efficient learning in a digital environment. In addition, these findings are

consistent with previous studies, which support the effectiveness of VR in disaster preparedness training (15-18).

The majority of participants noted the VR interface to be user-friendly and felt that the scenarios realistically simulated real-life disaster situations. A well-designed, user-friendly interface plays a crucial role in enhancing learning by fostering active engagement and improving knowledge retention (19). Furthermore, investing in user-friendly VR interfaces can enhance immersion and satisfaction in learning experiences, ultimately improving training effectiveness (20). Consistent with this, most participants reported that their actions in the VR environment would translate to real-world responses. Participants' feedback from DRM VR simulations provides valuable insights into their perceived effectiveness and acceptance. In a study evaluating VR as a training tool for public health Rapid Response Teams, participants reported high usability and excitement, indicating a positive reception of VR-based training methods (21).

Our study challenges the common belief that VR experiences in disaster response medical education led to a high incidence of cybersickness, despite previous research reporting discomfort rates ranging from 20% to 80% among users (8,22-23). Our findings indicate a more favorable level of acceptance. Only 30% of participants reported visual discomfort, and just 14.3% experienced dizziness following VR training. Several factors may explain these lower rates of cybersickness. One key factor is age. Kim H et al. (22) reported a higher prevalence of cybersickness in individuals aged 40 to 59 compared to those aged 19 to 39. Similarly, Keshavarz et al. (24) stated that older adults (65 and above) are more susceptible than younger individuals (18 to 39 years old). Another critical factor is exposure time. To minimize adverse effects, we limited VR sessions to 30 minutes. This is particularly relevant, as a significant incidence of cybersickness has been observed in participants after a 30-minute VR session (25). Moreover, our study incorporated VR stereo 3D glasses, which are recognized for being less disorienting and isolating (26). This feature may have contributed to the lower incidence of discomfort reported by our participants.

Performance assessments further demonstrated the effectiveness of VR training, with 60% of participants achieving a high level of proficiency. VR simulations surpass traditional training methods by providing a fully immersive and realistic learning environment (27). In contrast to conventional tabletop exercises, VR-based training immerses participants in authentic disaster scenarios, enhancing the development of preparation plans (28). Apart from that, in conventional field disaster training, students are typically assigned specific roles, such as triaging patients or delivering treatment at the CCS (29). As a result, they have limited exposure to the broader disaster management process, restricting their understanding of how various components function together. To address this gap, our module was designed

as a structured, step-by-step simulation of an out-of-hospital disaster response. The scenario begins with reporting to the on-scene commander, identifying an appropriate location for the CCS, and setting it up with the necessary equipment for both indoor and outdoor operations. Participants then triage victims based on injury severity, provide appropriate on-site management, and make transport decisions according to patient acuity levels. By fully immersing participants in this structured simulation, VR enhances their comprehension of disaster response coordination, decision making, and interprofessional collaboration (30). This comprehensive training approach likely contributed to the strong performance observed among participants in this study. By creating carefully designed and realistic scenarios, participants demonstrated strong proficiency, with an additional 40% achieving a moderate score.

LIMITATIONS

This study has several limitations. First, the use of written closed-ended questionnaires, while structured and practical, may have limited the immersive experience of the VR DRM training. To enhance fidelity in future research, incorporating voice prompts powered by large language models could offer a more interactive and engaging assessment, providing deeper insights without compromising the current study's validity. Second, the study population varied in terms of working experience and exposure to disaster situations. Considering their differing levels of prior training, establishing a uniform baseline of experience was challenging, which may have influenced their perceptions of the VR training. Lastly, the VR stereo 3D glasses were not well-fitted for participants who wore spectacles, potentially reducing their sense of immersion and making it harder for them to fully engage in the simulated disaster scenarios. Thus, future studies should consider optimizing VR equipment to enhance user comfort and immersion.

CONCLUSION

This study highlights VR's growing role in DRM education, emphasizing its usability, realism, and skill enhancement. With continued refinement, VR can bridge gaps in traditional training, offering an immersive approach to better prepare healthcare professionals. While concerns about cybersickness exist, by optimizing VR exposure time and using less disorienting hardware, the incidence of adverse effects can be minimized, thus making VR a viable and effective DRM training tool.

Acknowledgements

The authors would like to express their sincere gratitude to the staff of the Department of Emergency Medicine for their invaluable support and contributions to this study. Their dedication and assistance were instrumental in the successful completion of this research. This study was financially supported by Integrasi Erat Sdn Bhd under grant FF-2022-164.

REFERENCES

1. Siman-Tov M, Davidson B, Adini B. Maintaining preparedness to severe though infrequent threats—can it be done?. *International journal of environmental research and public health*. 2020 Apr;17(7):2385.
2. American College of Emergency Physicians. Role of emergency physicians in disaster preparedness and response: impact of COVID pandemic [Internet]. Irving, TX: American College of Emergency Physicians; [cited 2025 Feb 19]. Available from: <https://www.acep.org/patient-care/policy-statements/role-of-emergency-physicians-in-disaster-preparedness-and-response-impact-of-covid-pandemic>
3. Walsh L, Subbarao I, Gebbie K, Schor KW, Lyznicki J, Strauss-Riggs K, Cooper A, Hsu EB, King RV, Mitas II JA, Hick J. Core competencies for disaster medicine and public health. *Disaster medicine and public health preparedness*. 2012 Mar;6(1):44-52.
4. Hsu EB, Li Y, Bayram JD, Levinson D, Yang S, Monahan C. State of virtual reality based disaster preparedness and response training. *PLoS currents*. 2013 Apr 24;5.
5. Chapman K, Arbon P. Are nurses ready?: Disaster preparedness in the acute setting. *Australasian Emergency Nursing Journal*. 2008 Aug 1;11(3):135-44.

6. Sarin RR, Cattamanchi S, Alqahtani A, Aljohani M, Keim M, Ciottone GR. Disaster education: a survey study to analyze disaster medicine training in emergency medicine residency programs in the United States. *Prehospital and disaster medicine*. 2017 Aug;32(4):368-73.
7. Cohen D, Sevdalis N, Taylor D, Kerr K, Heys M, Willett K, Batrick N, Darzi A. Emergency preparedness in the 21st century: training and preparation modules in virtual environments. *Resuscitation*. 2013 Jan 1;84(1):78-84.
8. Farra SL, Miller ET, Hodgson E. Virtual reality disaster training: Translation to practice. *Nurse Education in Practice*. 2015 Jan 1;15(1):53-7.
9. Analyti E, Charitou R, Pesmatzoglou E, Stavrogiannopoulou M, Schoina I, Travlou C, Mitroyanni E. Virtual Reality in Education: Transforming Learning through Immersive Technology. *Technium Education and Humanities*. 2024 Oct 4;10:1-1.
10. Zhang H, Yu L, Ji M, Cui Y, Liu D, Li Y, Liu H, Wang Y. Investigating high school students' perceptions and presences under VR learning environment. In *Cross Reality (XR) and Immersive Learning Environments (ILEs) in Education 2023* Nov 14 (pp. 97-117). Routledge.
11. Andonova V, Reinoso-Carvalho F, Jimenez Ramirez MA, Carrasquilla D. Does multisensory stimulation with virtual reality (VR) and smell improve learning? An educational experience in recall and creativity. *Frontiers in Psychology*. 2023 Jun 15;14:1176697.
12. Yazdi M. Augmented Reality (AR) and Virtual Reality (VR) in Maintenance Training. In *Advances in Computational Mathematics for Industrial System Reliability and Maintainability 2024* Feb 25 (pp. 169-183). Cham: Springer Nature Switzerland.
13. Barney S, Khurum M, Petersen K, Unterkalmsteiner M, Jabangwe R. Improving students with rubric-based self-assessment and oral feedback. *IEEE transactions on Education*. 2011 Nov 4;55(3):319-25.

14. Hattie J, Timperley H. The power of feedback. *Review of educational research*. 2007 Mar;77(1):81-112.
15. Pucher PH, Batrick N, Taylor D, Chaudery M, Cohen D, Darzi A. Virtual-world hospital simulation for real-world disaster response: design and validation of a virtual reality simulator for mass casualty incident management. *Journal of trauma and acute care surgery*. 2014 Aug 1;77(2):315-21.
16. Bourhim EM, Cherkaoui A. Efficacy of virtual reality for studying people's pre-evacuation behavior under fire. *International Journal of Human-Computer Studies*. 2020 Oct 1;142:102484.
17. Alshowair A, Bail J, AlSuwailem F, Mostafa A, Abdel-Azeem A. Use of virtual reality exercises in disaster preparedness training: A scoping review. *SAGE Open Medicine*. 2024 Apr;12:20503121241241936.
18. Jung Y. Virtual reality simulation for disaster preparedness training in hospitals: integrated review. *Journal of Medical Internet Research*. 2022 Jan 28;24(1):e30600.
19. Ma CW, Cheng PS, Chan YS, Tipoe GL. Virtual reality: a technology to promote active learning of physiology for students across multiple disciplines. *Advances in Physiology Education*. 2023 Sep 1;47(3):594-603.
20. Serna-Mendiburu GM, Guerra-Tamez CR. Shaping the future of creative education: the transformative power of VR in art and design learning. *InFrontiers in Education* 2024 Apr 5 (Vol. 9, p. 1388483). Frontiers Media SA.
21. Bugli D, Dick L, Wingate KC, Driscoll S, Beck D, Walsh B, Greiner AL. Training the public health emergency response workforce: a mixed-methods approach to evaluating the virtual reality modality. *BMJ open*. 2023 May 1;13(5):e063527.

22. Kim H, Kim DJ, Chung WH, Park KA, Kim JD, Kim D, Kim K, Jeon HJ. Clinical predictors of cybersickness in virtual reality (VR) among highly stressed people. *Scientific reports*. 2021 Jun 9;11(1):12139.
23. Rebenitsch L, Owen C. Estimating cybersickness from virtual reality applications. *Virtual Reality*. 2021 Mar;25(1):165-74.
24. Keshavarz B, Ramkhalawansingh R, Haycock B, Shahab S, Campos JL. Comparing simulator sickness in younger and older adults during simulated driving under different multisensory conditions. *Transportation research part F: traffic psychology and behaviour*. 2018 Apr 1;54:47-62.
25. da Silva Marinho A, Terton U, Jones CM. Cybersickness and postural stability of first time VR users playing VR videogames. *Applied Ergonomics*. 2022 May 1;101:103698.
26. Steantycip. VR Glasses: Everything You Need to Know [Internet]. 2023 [cited 2025 Feb 19]. Available from: <https://steantycip.com/blog/vr-glasses/>
27. Liu JY, Yin YH, Kor PP, Cheung DS, Zhao IY, Wang S, Su JJ, Christensen M, Tyrovolas S, Leung AY. The effects of immersive virtual reality applications on enhancing the learning outcomes of undergraduate health care students: systematic review with meta-synthesis. *Journal of Medical Internet Research*. 2023 Mar 6;25:e39989.
28. Magi CE, Bambi S, Iovino P, El Aoufy K, Amato C, Balestri C, Rasero L, Longobucco Y. Virtual reality and augmented reality training in disaster medicine courses for students in nursing: a scoping review of adoptable tools. *Behavioral Sciences*. 2023 Jul 24;13(7):616.
29. Rajagopal AB, Jasperse N, Osborn MB. Simulated mass casualty incident triage exercise for training medical personnel. *Journal of Education & Teaching in Emergency Medicine*. 2020 Oct;5(4):SG1.

30. Al Thobaity A. Overcoming challenges in nursing disaster preparedness and response: an umbrella review. *BMC nursing*. 2024 Aug 14;23(1):562.

Early view